

THE VACUUM, LIGHT SPEED, AND THE REDSHIFT

PRELIMINARY NOTE

This document is intended to give an overview of the main conclusions reached from recent developments in light-speed research. In order to do this effectively, it has been necessary to include background information which, for a few, will already be well-known. However, for the sake of the majority who are not conversant with these areas of physics, it was felt important to include this information. While this overview is comprehensive, the actual derivation of many conclusions is beyond its scope. These derivations have, nevertheless, been fully performed in a major scientific paper using standard maths and physics coupled with observational data. Full justification of the conclusions mentioned here can be found in that thesis. Currently, that paper in which the new model is presented, is being finalised for peer review and will be made available once this whole process is complete.

THE VACUUM

During the 20th century, our knowledge regarding space and the properties of the vacuum has taken a considerable leap forward. The vacuum is more unusual than many people realise. It is popularly considered to be a void, an emptiness, or just 'nothingness.' This is the definition of a *bare vacuum* [1]. However, as science has learned more about the properties of space, a new and contrasting description has arisen, which physicists call the *physical vacuum* [1].

To understand the difference between these two definitions, imagine you have a perfectly sealed container. First remove all solids and liquids from it, and then pump out all gases so no atoms or molecules remain. There is now a vacuum in the container. It was this concept in the 17th century that gave rise to the definition of a *vacuum* as a totally empty volume of space. It was later discovered that, although this vacuum would not transmit sound, it would transmit light and all other wavelengths of the electromagnetic spectrum. Starting from the high energy side, these wavelengths range from very short wavelength gamma rays, X-rays, and ultra-violet light, through the rainbow spectrum of visible light, to low energy longer wavelengths including infra-red light, microwaves and radio waves.

THE ENERGY IN THE VACUUM

Then, late in the 19th century, it was realised that the vacuum could still contain heat or thermal radiation. If our container with the vacuum is now perfectly insulated so no heat can get in or out, and if it is then cooled to absolute zero, all thermal radiation will have been removed. Does a complete vacuum now exist within the container? Surprisingly, this is not the case. Both theory and experiment show that this vacuum still contains measurable energy. This energy is called the *zero-point energy* (ZPE) because it exists even at absolute zero.

The ZPE was discovered to be a universal phenomenon, uniform and all-pervasive on a large scale. Therefore, its existence was not suspected until the early 20th century. In 1911, while working with a series of equations describing the behaviour of radiant energy from a hot body, Max Planck found that the observations required a term in his equations that did not depend on temperature. Other physicists, including Einstein, found similar terms appearing in their own equations. The implication was that, even at absolute zero, each body will have some residual energy. Experimental evidence soon built up hinting at the existence of the ZPE, although its fluctuations do not become significant enough to be observed until the atomic level is attained. For example [2], the ZPE can explain why cooling alone will never freeze liquid helium. Unless pressure is applied, these ZPE fluctuations prevent helium's atoms from getting close enough to permit solidification. In electronic circuits another problem surfaces because ZPE fluctuations cause a random "noise" that places limits on the level to which signals can be amplified.

The magnitude of the ZPE is truly large. It is usually quoted in terms of energy per unit of volume which is referred to as **energy density**. Well-known physicist Richard Feynman and others [3] have pointed out that the amount of ZPE in one cubic centimetre of the vacuum "is greater than the energy density in an atomic nucleus" [4]. Indeed, it has been stated that [5]: "Formally, physicists attribute an infinite amount of energy to this background. But, even when they impose appropriate cutoffs at high frequency, they estimate conservatively that the zero-point density is comparable to the energy density inside an atomic nucleus." In an atomic nucleus alone, the energy density is of the order of 10^{44} ergs per cubic centimetre. (An **erg** is defined as "the energy expended or work done when a mass of 1 gram undergoes an acceleration of 1 centimetre per second per second over a distance of 1 centimetre.")

Estimates of the energy density of the ZPE therefore range from at least 10^{44} ergs per cubic centimetre up to infinity. For example, Jon Noring made the statement that "Quantum Mechanics predicts the energy density [of the ZPE] is on the order of an incomprehensible 10^{98} ergs per cubic centimetre." Prigogine and Stengers also analysed the situation and provided estimates of the size of the ZPE ranging from 10^{100} ergs per cubic centimetre up to infinity. In case this is dismissed as fanciful, Stephen M. Barnett from the University of Oxford, writing in *Nature* (March 22, 1990, p.289), stated: "The mysterious nature of the vacuum [is] revealed by quantum electrodynamics. It is not an empty nothing, but contains randomly fluctuating electromagnetic fields...with an infinite zero-point energy." In actual practice, recent work suggests there may be an upper limit for the estimation of the ZPE at about 10^{114} ergs per cubic centimetre (this upper limit is imposed by the Planck length, as discussed below).

In order to appreciate the magnitude of the ZPE in each cubic centimetre of space, consider a conservative estimate of 10^{52} ergs/cc. Most people are familiar with the light bulbs with which we illuminate our houses. The one in my office is labeled as 150 watts. (A **watt** is defined as 10^7 ergs per second.) By comparison, our sun radiates energy at the rate of 3.8×10^{20} watts. In our galaxy there are in excess of 100 billion stars. If we assume they all radiate at about the same intensity as our sun, then the amount of energy expended by our entire galaxy of stars shining for one million years is roughly equivalent to the energy locked up in one cubic centimetre of space.

THE "GRANULAR STRUCTURE" OF SPACE

In addition to the ZPE, there is another aspect of the physical vacuum that needs to be presented. When dealing with the vacuum, size considerations are all-important. On a large scale the physical vacuum has properties that are uniform throughout the cosmos, and seemingly smooth and featureless. However, on an atomic scale, the vacuum has been described as a "seething sea of activity" [2], or "the seething vacuum" [5]. It is in this realm of the very small that our understanding of the vacuum has increased. The size of the atom is about 10^{-8} centimetres. The size of an atomic particle, such as an electron, is about 10^{-13} centimetres. As the scale becomes smaller, there is a major change at the **Planck length** (1.616×10^{-33} centimetres), which we will designate as L^* [6]. In 1983, F. M. Pipkin and R. C. Ritter pointed out in *Science* (vol. 219, p.4587), that "the Planck length is a length at which the smoothness of space breaks down, and space assumes a granular structure."

This "granular structure" of space, to use Pipkin and Ritter's phrase, is considered to be made up of Planck particles whose diameter is equal to L^* , and whose mass is equal to a fundamental unit called the **Planck mass**, M^* , (2.177×10^{-5} grams). These Planck particles form the basis for various cosmological theories such as strings, super strings, 10-dimensional space, and so on. During the last hundred years, physicists have discovered that atomic particles such as electrons or protons, have a wave-form associated with them. This is termed the **wave/particle duality of matter**. These waves are called **deBroglie** waves and vary inversely with mass [7]. That is to say, the heavier the particle, the shorter its wavelength. This means that because a proton is more massive, its wavelength is shorter than an electron's. What is interesting is that Planck particles have a diameter L^* that is equal to their deBroglie wavelength.

The physical vacuum of space therefore appears to be made up of an all-pervasive sea of Planck particles whose density is an unbelievable 3.6×10^{93} grams per cubic centimetre. It might be wondered how anything can move through such a medium. It is because deBroglie wavelengths of elementary particles are so long compared with the Planck length, L^* , that the vacuum is 'transparent' to these elementary particles.

It is for the same reason that long wavelength infra-red light can travel through a dense cloud in space and reveal what is within instead of being absorbed, and why light can pass through dense glass. Therefore, motion of elementary particles through the vacuum will be effortless, as long as these particles do not have energies of the magnitude of what is referred to as **Planck energy**, or $M \cdot c^2$ ('c' is the velocity of light). Atomic particles of that energy would simply be absorbed by the structure of the vacuum. From the figures for the density given above, the energy associated with this Planck particle sea making up the physical vacuum can be calculated to be of the order of 10^{14} ergs per cubic centimetre, the same as the maximum value for the ZPE.

TWO THEORIES DESCRIBING THE VACUUM

Currently, there are two theories that describe the behaviour and characteristics of the physical vacuum and the ZPE at the atomic or sub-atomic level: the *quantum electro-dynamic* (QED) model [8], and the somewhat more recent *stochastic electro-dynamic* (SED) model [9,10]. They both give the same answers mathematically, so the choice between them is one of aesthetics. In some cases the QED model gives results that are easier to visualise; in other cases the SED model is better. Importantly, both come to the same conclusion that even at absolute zero the physical vacuum has an inherent energy density. The origin of this energy is discussed later. For now, the focus of attention is on the observable effects of this energy. The QED model maintains that the zero-point energy reveals its existence through the effects of sub-atomic virtual particles. By contrast, the SED approach affirms that the ZPE exists as electromagnetic fields or waves whose effects explain the observed phenomena equally well. Let us look at both in a little more detail.

THE QED MODEL OF THE VACUUM

At the atomic level, the QED model proposes that, because of the high inherent energy density within the vacuum, some of this energy can be temporarily converted to mass. This is possible since energy and mass can be converted from one to the other according to Einstein's famous equation [$E = mc^2$], where 'E' is energy, 'm' is mass, and 'c' is the speed of light. On this basis, the QED model proposes that the ZPE permits short-lived particle/antiparticle pairs (such as a positive and negative pion, or perhaps an electron and positron) to form and almost immediately annihilate each other [2,11]. These particle/antiparticle pairs are called *virtual particles*. Virtual particles are distinct from Planck particles which make up the structure of the vacuum. While virtual particles are, perhaps, about 10^{-13} centimetres diameter, Planck particles are dramatically smaller at about 10^{-33} cm. Virtual particles wink in and out of existence incredibly quickly. The exact relationship between the energy of these particles and the brief time of their existence is explained in quantum theory by *Heisenberg's uncertainty principle*.

The Heisenberg uncertainty principle states that the uncertainty of time multiplied by the uncertainty of the energy is closely approximated to *Planck's constant* 'h' divided by 2π , (π stands for the mathematical symbol pi in the standard relation $p = \pi = 3.14159\dots$). This quantum uncertainty, or indeterminacy, governed by the value of 'h', imposes fundamental limitations on the precision with which a number of physical quantities associated with atomic processes can be measured. In the case under consideration here, the uncertainty principle permits these virtual particle events to occur as long as they are completed within an extraordinarily brief period of time, which is of the order of 10^{-23} seconds [5]. According to this QED model, an atomic particle such as a proton or electron, even when entirely alone in a vacuum at absolute zero, is continually emitting and absorbing these virtual particles from the vacuum [12].

Consequently, a proton or electron is considered to be the centre of constant activity; it is surrounded by a cloud of virtual particles with which it is interacting [12]. In the case of the electron, physicists have been able to penetrate a considerable way into this virtual particle cloud. They have found that the further into the cloud they go, the smaller, more compact and point-like the electron becomes. At the same time they have discovered there is a more pronounced negative charge associated with the electron the further they penetrated into this cloud [13]. These virtual particles act in such a way as to screen the full electronic charge. There is a further important effect verified by observation and experiment: the absorption and emission of these virtual particles also causes the electron's "jitter motion" in a vacuum at absolute zero.

As such, this jittering, or *Zitterbewegung*, as it is officially called [14], constitutes evidence for the existence of virtual particles and the ZPE of the vacuum.

THE SED MODEL OF THE VACUUM

In the SED approach, the vacuum at the atomic or sub-atomic level may be considered to be inherently comprised of a turbulent sea of randomly fluctuating electro-magnetic fields or waves. These waves exist at all wavelengths longer than the Planck length L^* . At the macroscopic level, these all-pervasive *zero-point fields* (ZPF) are homogeneous and isotropic, which means they have the same properties uniformly in every direction throughout the whole cosmos. Furthermore, observation shows that this *zero-point radiation* (ZPR) must be “*Lorentz invariant*” [1]. This means that it must look the same to two observers no matter what the velocity of these observers is with respect to each other. Note that this Lorentz invariance makes the ZPF crucially different from any of the 19th century concepts of an ether [15]. The old ether concept indicated absolute velocity through the ether could be determined. However, the Lorentz invariant condition indicates that the zero-point radiation will look the same to all observers regardless of their relative velocities.

Importantly, with the SED approach, *Planck’s quantum constant*, ‘ h ’, becomes a measure of the strength of the ZPF. This situation arises because the fluctuations of the ZPF provide an irreducible random noise at the atomic level that is interpreted as the innate uncertainty described by Heisenberg’s uncertainty principle [4,16]. Therefore, the zero-point fields are the ultimate source of this fundamental limitation with which we can measure some atomic phenomena and, as such, give rise to the indeterminacy or uncertainty of quantum theory mentioned above. In fact, Nelson pointed out in 1966 that if the ZPR had been discovered at the beginning of the 20th century, then classical mechanics plus the ZPR could have formulated nearly all the results developed by quantum mechanics [17, 4].

In the SED explanation, the *Zitterbewegung* is accounted for by the random fluctuations of the ZPF, or waves, as they impact upon the electron and jiggle it around. There is also evidence for the existence of the zero-point energy in this model by something called the surface *Casimir effect*, predicted Hendrik Casimir, the Dutch scientist, in 1948 and confirmed nine years later by M. J. Sparnaay of the Philips Laboratory in Eindhoven, Holland [1]. The Casimir effect can be demonstrated by bringing two large metal plates very close together in a vacuum. When they are close, but not touching, there is a small but measurable force that pushes them together. The SED theory explains this simply. As the metal plates get closer, they end up excluding all wavelengths of the ZPF between the plates except the very short ones that are a sub-multiple of the plates’ distance apart. In other words, all the long wavelengths of the ZPF are now acting on the plates from the outside. The combined radiation pressure of these external waves then forces the plates together [5,16]. The same effect can be seen on the ocean. Sailors have noted that if the distance between two boats is less than the distance between two wave crests (or one wavelength), the boats are forced towards each other.

The Casimir effect is directly proportional to the area of the plates. However, unlike other possible forces with which it may be confused, the Casimir force is inversely proportional to the fourth power of the plates’ distance apart [18]. For plates with an area of one square centimetre separated by 0.5 thousandths of a millimetre, this force is equivalent to a weight of 0.2 milligrams. In January of 1997, Steven Lamoreaux reported verification of these details by an experiment reported in *Physical Review Letters* (vol.78, p5).

The surface Casimir effect therefore demonstrates the existence of the ZPE in the form of electromagnetic waves. Interestingly, Haisch, Rueda, Puthoff and others point out that there is a microscopic version of the same phenomenon. In the case of closely spaced atoms or molecules the all-pervasive ZPF result in short-range attractive forces that are known as *van der Waals forces* [4,16]. It is these attractive forces that permit *real gases* to be turned into liquids [2]. (When an ‘ideal’ gas is compressed, it behaves in a precise way. When a real gas is compressed, its behaviour deviates from the ideal equation [19]).

The common objections to the actual existence of the zero-point energy centre around the idea that it is simply a theoretical construct. However the presence of both the Casimir effect and the *Zitterbewegung*, among other observational evidences, prove the reality of the ZPE.

LIGHT AND THE PROPERTIES OF SPACE

This intrinsic energy, the ZPE, which is inherent in the vacuum, gives free space its various properties. For example, the magnetic property of free space is called the *permeability* while the corresponding electric property is called the *permittivity*. Both of these are affected uniformly by the ZPE [20]. If they were not, the electric and magnetic fields in travelling light waves would no longer bear a constant ratio to each other, and light from distant objects would be noticeably affected [21]. Since the vacuum permeability and permittivity are also energy-related quantities, they are directly proportional to the energy per unit volume (the energy density) of the ZPE [20]. It follows that if the energy density of the ZPE ever increased, then there would be a proportional increase in the value of both the permeability and permittivity.

Because light waves are an electro-magnetic phenomenon, their motion through space is affected by the electric and magnetic properties of the vacuum, namely the permittivity and permeability. To examine this in more detail we closely follow a statement by Lehrman and Swartz [22]. They pointed out that light waves consist of changing electric fields that generate changing magnetic fields. This then regenerates the electric field, and so on. The wave travels by transferring energy from the electric field to the magnetic field and back again. The magnetic field resulting from the change in the electric field must be such as to oppose the change in the electric field, according to *Lenz's Law*. This means that the magnetic property of space has a kind of inertial property inhibiting the rapid change of the fields. The magnitude of this property is the *magnetic constant* of free space 'U' which is usually called the *magnetic permeability* of the vacuum.

The electric constant, or permittivity, of free space is also important, and is related to electric charges. A charge represents a kind of electrical distortion of space, which produces a force on neighbouring charges. The constant of proportionality between the interacting charges is $1/Q$, which describes a kind of electric elastic property of space. The quantity Q is usually called the *electric permittivity* of the vacuum. It is established physics that the velocity of a wave motion squared is proportional to the ratio of the elasticity over the inertia of the medium in which it is travelling. In the case of the vacuum and the speed of light, c , this standard equation becomes

$$c^2 = 1 / (UQ)$$

As noted above, both U and Q are directly proportional to the energy density of the ZPE. It therefore follows that any increase in the energy density of the ZPF will not only result in a proportional increase in U and Q , but will also cause a decrease in the speed of light, c .

WHY ATOMS DON'T SELF-DESTRUCT

But it is not only light that is affected by these properties of the vacuum. It has also been shown that the atomic building blocks of matter are dependent upon the ZPE for their very existence. This was clearly demonstrated by Dr. Hal Puthoff of the Institute for Advanced Studies in Austin, Texas. In *Physical Review D*, vol. 35:10, and later in *New Scientist* (28 July 1990), Puthoff started by pointing out an anomaly. According to classical concepts, an electron in orbit around a proton should be radiating energy. As a consequence, as it loses energy, it should spiral into the atomic nucleus, causing the whole structure to disappear in a flash of light. But that does not happen. When you ask a physicist why it does not happen, you will be told it is because of *Bohr's quantum condition*. This quantum condition states that electrons in specific orbits around the nucleus do **not** radiate energy. But if you ask why not, or alternatively, if you ask why the classical laws of electro-magnetics are violated in this way, the reply is generally vague and less than satisfactory [4].

Instead of ignoring the known laws of physics, Puthoff approached this problem with the assumption that the classical laws of electro-magnetics were valid, and that the electron is therefore losing energy as it speeds in its orbit around the nucleus. He also accepted the experimental evidence for the existence of the ZPE in the form of randomly fluctuating electro-magnetic fields or waves. He calculated the power the electron lost as it moved in its orbit, and then calculated the power that the electron gained from the ZPF. The two turned out to be identical; the loss was exactly made up for by the gain. It was like a child on a

swing: just as the swing started to slow, it was given another push to keep it going. Puthoff then concluded that without the ZPF inherent within the vacuum, every atom in the universe would undergo instantaneous collapse [4, 23]. In other words, the ZPE is maintaining all atomic structures throughout the entire cosmos.

THE RAINBOW SPECTRUM

Knowing that light itself is affected by the zero-point energy, phenomena associated with light need to be examined. When light from the sun is passed through a prism, it is split up into a spectrum of seven colours. Falling rain acts the same way, and the resulting spectrum is called a rainbow. Just like the sun and other stars making up our own galaxy, distant galaxies each have a rainbow spectrum. From 1912 to 1922, Vesto Slipher at the Lowell Observatory in Arizona recorded accurate spectrographic measurements of light from 42 galaxies [24, 25]. When an electron drops from an outer atomic orbit to an inner orbit, it gives up its excess energy as a flash of light of a very specific wavelength. This causes a bright emission line in the colour spectrum.

However when an electron jumps to a higher orbit, energy is absorbed and instead of a bright emission line, the reverse happens – a dark absorption line appears in the spectrum. Each element has a very specific set of spectral lines associated with it. Within the spectra of the sun, stars or distant galaxies these same spectral lines appear.

THE REDSHIFT OF LIGHT FROM GALAXIES

Slipher noted that in distant galaxies this familiar pattern of lines was shifted systematically towards the red end of the spectrum. He concluded that this redshift of light from these galaxies was a *Doppler effect* caused by these galaxies moving away from us. The Doppler effect can be explained by what happens to the pitch of a siren on a police car as it moves away from you. The tone drops Slipher concluded that the redshift of the spectral lines to longer wavelengths was similarly due to the galaxies receding from us. For that reason, this redshift is usually expressed as a velocity, even though as late as 1960 some astronomers were seeking other explanations [25]. In 1929, Edwin Hubble plotted the most recent distance measurements of these galaxies on one axis, with their redshift recession velocity on the other. He noted that the further away the galaxies were, the higher were their redshifts [24].

It was concluded that if the redshift represented receding galaxies, and the redshift increased in direct proportion to the galaxies distances from us, then the entire universe must be expanding [24]. The situation is likened to dots on the surface of a balloon being inflated. As the balloon expands, each dot appears to recede from every other dot. A slightly more complete picture was given by relativity theory. Here space itself is considered to be expanding, carrying the galaxies with it. According this interpretation, light from distant objects has its wavelength stretched or reddened in transit because the space in which it is travelling is expanding.

THE REDSHIFT GOES IN JUMPS

This interpretation of the redshift is held by a majority of astronomers. However, in 1976, William Tift of the Steward Observatory in Tucson, Arizona, published the first of a number of papers analyzing redshift measurements. He observed that the redshift measurements did not change smoothly as distance increased, but went in jumps: in other words they were *quantised* [26]. Between successive jumps, the redshift remained fixed at the value it attained at the last jump. This first study was by no means exhaustive, so Tift investigated further. As he did so, he discovered that the original observations that suggested a quantised redshift were strongly supported wherever he looked [27 - 34]. In 1981 the extensive Fisher-Tully redshift survey was completed. Because redshift values in this survey were not clustered in the way Tift had noted earlier, it looked as if redshift quantisation could be ruled out. However, in 1984 Tift and Cocke pointed out that the motion of the sun and its solar system through space produces a genuine Doppler effect of its own, which adds or subtracts a little to every redshift measurement. When this true Doppler effect was subtracted from all the observed redshifts, it produced strong evidence for the quantisation of redshifts across the entire sky [35, 36].

The initial quantisation value that Tifft discovered was a redshift of 72.46 kilometres per second in the Coma cluster of galaxies. Subsequently it was discovered that quantisation figures of up to 13 multiples of 72.46 km/s existed. Later work established a smaller quantisation figure just half of this, namely 36.2 km/s. This was subsequently supported by Guthrie and Napier who concluded that 37.6 km/s was a more basic figure, with an error of 2 km/s [37-39]. After further observations, Tifft announced in 1991 that these and other redshift quantisations recorded earlier were simply higher multiples of a basic quantisation figure [40]. That figure turned out to be 8.05 km/s, which when multiplied by 9 gave the original 72.46 km/s value. Alternatively, when 8.05 km/s is multiplied by 9/2 the 36.2 km/s result is obtained. However, Tifft noted that this 8.05 km/s was not in itself the most basic result as observations revealed a 8.05/3 km/s, or 2.68 km/s, quantisation, which was even more fundamental [40]. Accepting this result at face value suggests that the redshift is quantised in fundamental steps of 2.68 km/s across the cosmos.

RE-EXAMINING THE REDSHIFT

If redshifts were truly a result of an expanding universe, the measurements would be smoothly distributed, showing all values within the range measured. This is the sort of thing we see on a highway, with cars going many different speeds within the normal range of driving speeds. However the redshift, being quantised, is more like the idea of those cars each going in multiples of, say, 5 kilometres an hour. Cars don't do that, but the redshift does. This would seem to indicate that something other than the expansion of the universe is responsible for these results.

We need to undertake a re-examination of what is actually being observed in order to find a solution to the problem. It is this solution to the redshift problem that introduces a new cosmological model. In this model, atomic behaviour and light-speed throughout the cosmos are linked with the ZPE and properties of the vacuum.

The prime definition of the redshift, 'z', involves two measured quantities. They comprise the observed change in wavelength 'D' of a given spectral line when compared with the laboratory standard 'W'. The ratio of these quantities [$D/W = z$] is a dimensionless number that measures the redshift [41]. However, it is customarily converted to a velocity by multiplying it by the current speed of light, 'c' [41]. The redshift so defined is then 'cz', and it is this cz which is changing in steps of 2.68 km/s. Since the laboratory standard wavelength 'W' is unaltered, it then follows that as [$z = D/W$] is systematically increasing in discrete jumps with distance, then D must be increasing in discrete jumps also. Now D is the difference between the observed wavelength of a given spectral line and the laboratory standard [41]. This suggests that emitted wavelengths are becoming longer in quantum jumps with increasing distance (or with look-back time). During the time between jumps, the emitted wavelengths remain unchanged from the value attained at the last jump.

The basic observations therefore indicate that the wavelengths of all atomic spectral lines have changed in discrete jumps throughout the cosmos with time. This could imply that all atomic emitters within each galaxy may be responsible for the quantised redshift, rather than the recession of those galaxies or universal expansion. Importantly, the wavelengths of light emitted from atoms are entirely dependent upon the energy of each atomic orbit. According to this new way of interpreting the data, the redshift observations might indicate that the energy of every atomic orbit in the cosmos simultaneously undergoes a series of discrete jumps with time. How could this be possible?

ATOMIC ORBITS AND THE REDSHIFT

The explanation may well be found in the work of Hal Puthoff. Since the ZPE is sustaining every atom and maintaining the electrons in their orbits, it would then also be directly responsible for the energy of each atomic orbit. In view of this, it can be postulated that if the ZPE were lower in the past, then these orbital energies would probably be less as well. Therefore emitted wavelengths would be longer, and hence redder. Because the energy of atomic orbits is quantised or goes in steps [42], it may well be that any increase in atomic orbital energy can similarly only go in discrete steps. Between these steps atomic orbit energies would remain fixed at the value attained at the last step. In fact, this is the precise effect that Tifft's redshift data reveals.

The outcome of this is that atomic orbits would be unable to access energy from the smoothly increasing ZPF until a complete unit of additional energy became available. Thus, between quantum jumps all atomic processes proceed on the basis of energy conservation, operating within the framework of energy provided at the last quantum jump. Increasing energy from the ZPE will not affect the atom until a particular threshold is reached, at which time all the atoms in the universe react simultaneously.

THE SIZE OF THE ELECTRON

This new approach can be analysed further. Mathematically it is known that the strength of the electronic charge is one of several factors governing the orbital energies within the atom [42]. Therefore, for the orbital energy to change, a simultaneous change in the value of the charge of both the electron and the proton would be expected. Although we will only consider the electron here, the same argument holds for the proton as well.

Theoretically, the size of the electron should appear to increase at each quantum jump, becoming “larger” with time. The so-called *Compton radius* of the electron is 3.86×10^{-11} centimetres which, in the SED approach, is significant. Haisch, Rueda, and Puthoff [16] have pointed out that “*one defensible interpretation is that the electron really is a point-like entity, smeared out to its quantum dimensions by the ZPF fluctuations.*” This “*smearing out*” of the electronic charge by the ZPF involves vacuum polarisation and the *Zitterbewegung*. When the calculations are done in SED using these phenomena, the Compton radius for the electron is indeed obtained [16].

THE ELECTRONIC CHARGE

With this in mind, it might be anticipated, on the SED approach, that if the energy density of the ZPF increased, the “*point-like entity*” of the electron would be “*smeared out*” even more, thus appearing larger. This would follow since the *Zitterbewegung* would be more energetic, and vacuum polarization around charges would be more extensive. In other words, the electron’s apparent radius could increase at the quantum jump. But the formula for the *classical radius* of the electron takes it a step further. It links the electron radius with the electronic charge. A larger radius means a stronger charge, other factors being equal. Therefore, at the quantum jump, when a full quantum of additional energy becomes available to the atom from the ZPE, the electron’s radius might be expected to expand which, according to the formula, might indicate that the intrinsic charge had increased. The importance of this is that a greater electronic charge will result in a greater orbital energy, which means that wavelengths emitted by the atom will be shifted towards the blue end of the spectrum.

The QED model can explain this formula another way. There is a cloud of virtual particles around the “bare” electron interacting with it. When a full quantum increase in the vacuum energy density occurs, the strength of the charge increases. With a higher charge for the “*point-like entity*” of the electron, it would be expected that the size of the particle cloud would increase because of stronger vacuum polarisation and a more energetic *Zitterbewegung*. (Note that *vacuum polarisation* occurs because of a tendency for virtual particles to be attracted to charges of the opposite sign, while those of the same sign remain more distant [18, 43]). This larger cloud of virtual particles intimately associated with the “bare” electron would give rise to an increase in the perceived radius of the “dressed” electron since it also includes the particle cloud. In fact this “dressed” electron is the entity that has been observed classically, and the one to which both the *Compton radius* and *classical radius* formulae apply. This inevitably means that the virtual particle cloud partially screens the full value of the “bare” charge. Some experiments have probed deep into the virtual particle cloud and found the charge does indeed increase with penetration. In fact, the full value of the “bare” charge has yet to be determined [13, 44].

THE BOHR ATOM

Let us now be more specific about this new approach to orbit energies and their association with the redshift. The *Bohr model* of the atom has electrons going around the atomic nucleus in miniature orbits, like planets around the sun. Although more sophisticated models of the atom now exist, it has been

acknowledged in the past that the Bohr theory “*is still often employed as a first approximation*” [45 - 47]. Similarly, much of the recent work done on the ZPE and atoms in the SED approach has also been at Bohr theory level [23]. It has been stated that the motive has been to gain “*intuitive insights and calculational ease*” [16]. Accordingly, that approach is retained here.

In the Bohr model of the atom, two equations describe orbital energy [42]. In 1913, Niels Bohr quantised the first of these, the angular momentum equation. The **angular momentum** of an orbit is described mathematically by ‘ mvr ’, where ‘ m ’ is the mass of the electron, ‘ v ’ is its velocity in an orbit whose radius is ‘ r ’. Bohr pointed out that a close approximation to observed atomic behaviour is obtained if electrons are theoretically restricted to those orbits whose angular momentum is an integral multiple of $h / (2\pi)$; ($p = \pi$). Mathematically, that is written as

$$mvr = nh / (2\pi)$$

where ‘ n ’ is a whole number such as 1, 2, 3, etc., and is called the **quantum number**. As mentioned above, ‘ h ’ is **Planck’s quantum constant**. This procedure effectively describes a series of permitted orbits for electrons in any given atom. In so doing it establishes the spectral line structure for any specific atom. That much is standard physics. The new approach maintains the integrity of Bohr’s first equation, so at the instant of any quantum jump in orbital energy, the angular momentum would be conserved.

BOHR’S SECOND EQUATION

Bohr’s second equation describes the kinetic energy of the electron in an orbit of radius ‘ r ’. **Kinetic energy** is defined as $mv^2/2$. The standard equation for the kinetic energy of the orbit closest to the nucleus reads

$$mv^2 / 2 = e^2 / [(8\pi)Qr]$$

where ‘ e ’ is the charge on the electron, and ‘ Q ’ is the permittivity of the vacuum. This kinetic energy is equal in magnitude to the total energy of that closest orbit. When an electron falls from immediately outside the atom into that orbit, this energy is released as a photon of light. The energy ‘ E ’ of this photon has a wavelength ‘ W ’ and both of these terms are linked by the standard equation

$$E = hc / W$$

As shown later, observational evidence reveals the ‘ hc ’ component in this equation is an absolute constant at all times. The kinetic energy and the photon energy are thus equal. Again, this much is standard physics [42]. Accordingly, we can write the following equality from Bohr’s second equation:

$$mv^2/2 = e^2 / [(8\pi)Qr] = hc / W$$

A NEW QUANTUM CONDITION

If we now follow the lead of Bohr, and quantise his second equation, a solution to several difficulties is found. Since the redshift indicates that the wavelengths W of light emitted from the galaxies are quantised, let us then quantise wavelengths such that

$$W = nR^*$$

Here, the term ‘ n ’ is the **new quantum integer** which fulfils the same function as Bohr’s quantum number ‘ n ’. Furthermore, Planck’s quantum constant ‘ h ’ finds its parallel in ‘ R^* ’ which could be called the **Rydberg quantum wavelength** defined as

$$R^* = 8.3041 \times 10^{-11} \text{ centimetres}$$

‘ R^* ’ turns out to be intrinsically related to the **Rydberg constant** ‘ R ’, which links emitted wavelengths with atomic orbit energy [42]. This link was discovered by Johannes Robert Rydberg of Sweden in 1890. In fact,

over a century later, this model indicates that he had discovered more than he is being credited with. To see this, note the inverse of the Rydberg constant gives the Rydberg wavelength 'K' as

$$1 / R = K = 9.1127 \times 10^{-6} \text{ centimetres}$$

Also defined here is a dimensionless number, the *Rydberg number* 'R' such that

$$1 / \underline{R} = 9.1127 \times 10^{-6}$$

The Rydberg quantum wavelength is then defined as

$$R^* = 1 / (R\underline{R}) = K / \underline{R} = 8.3041 \times 10^{-11} \text{ centimetres}$$

This new quantisation procedure means that as 'n' decreases with time, the energy of the first Bohr orbit will increase in steps of

$$hc / (\underline{n}R^*) = mv^2 / 2 = e^2 / [(8p)Qr]$$

This holds because of two factors. First, if 'n' decreases with time, it will mimic the behaviour of the redshift which also decreases with time. High redshift values from distant objects necessarily mean high values for 'n' as well. Second, all orbit radii 'r' can be shown to remain unchanged throughout any quantum changes. If they were not, the abrupt change of size of every atom at the quantum jump would cause obvious flaws in crystals, which would be especially noticeable in ancient rocks. This new quantisation procedure effectively allows every atom in the cosmos to simultaneously acquire a new higher energy state for each of its orbits in proportion as the ZPE increases with time. In so doing, it opens the way for a solution to the redshift problem.

A QUANTUM REDSHIFT

In the Bohr atom, all orbit energies are scaled according to the energy of the orbit closest to the nucleus known as the *ground state orbit*. Therefore, if the ground state orbit has an energy change, all other orbits will scale their energy proportionally. This also means that wavelengths of emitted light will be scaled in proportion to the energy of the ground state orbit of the atom. Accordingly, if W_0 is any arbitrary emitted wavelength and W_1 is the wavelength of the ground state orbit, then the wavelength change at the quantum jump is given by

$$D = \underline{n}R^*W_0 / W_1$$

Now the redshift is defined as the change in wavelength D divided by the reference wavelength W. For the purposes of illustration, let us take the reference wavelength to be equal to that emitted when an electron falls into the ground state orbit for hydrogen. This wavelength is close to 9.12×10^{-6} centimetres. For this orbit, the value of D from the above equation is given by 8.304×10^{-11} centimetres. Therefore, the redshift

$$z = D / W = 9.105 \times 10^{-6}$$

and so the velocity change

$$cz = 2.73 \text{ km/sec}$$

This compares very favourably with Tiff's value of 2.68 km/sec for the quantum jumps in the redshift.

Imposing a quantum condition on the second Bohr equation for the atom therefore produces quantum changes in orbit energies and emitted wavelengths that accord with the observational evidence. This result also implies the quantised redshift may not be an indicator of universal expansion. Rather, this new model suggests it may be evidence that the ZPE has increased with time allowing atomic orbits to take up successively higher energy states.

AN INCREASING VACUUM ENERGY?

The key question then becomes, why should the ZPE increase with time? One basic tenet of the Big Bang and some other cosmologies is an initial rapid expansion of the universe. That initial rapid expansion is accepted here. However, the redshift can no longer be used as evidence that this initial expansion has continued until the present. Indeed, if space were continuing its uniform expansion, the precise quantisation of spectral line shifts that Tift has noted would be smeared out and lost. The same argument applies to cosmological contraction. This suggests that the initial expansion halted before redshifted spectral lines were emitted by the most distant galaxies, and that since then the universe has been static. In 1993, Jayant Narliker and Halton Arp published a paper in *Astrophysical Journal* (vol. 405, p. 51) which revealed that a static cosmos which contained matter was indeed stable against collapse.

However, the initial expansion was very important. As Paul S. Wesson [48], Martin Harwit [49] and others have shown, the physical vacuum initially acquired a potential energy in the form of an elasticity, tension, or stress as a result of the inflationary expansion of the cosmos. This might be considered to be akin to the tension, stress, or elasticity in the fabric of a balloon that has been inflated. Over time, this tensional energy changes its form. In exactly the same way that energy is liberated when liquid water changes to ice, so also the tensional energy of the vacuum is liberated in the form of radiation [50]. As Harwit points out, the energy residing in the elasticity of the vacuum (a form of potential energy) becomes converted into radiation (a form of kinetic energy) [49]. In the new model under consideration here, it is maintained that this potential energy becomes converted specifically into the zero-point radiation (ZPR) as the initial tension of the inflated cosmos ‘relaxes’ over time.

What is being proposed on this new model is that the ZPR content of the vacuum was low initially, but has built up with time as the potential energy of the elastic tension changed its form into the ZPE of the vacuum electro-magnetic fields. The redshift data indicate that this conversion of the vacuum elasticity into the ZPE essentially followed an exponential decay.

RECONSIDERING LIGHT-SPEED

It is at this point in the discussion that a consideration of light-speed becomes important. It has already been mentioned that an increase in vacuum energy density will result in an increase in the electrical permittivity and the magnetic permeability of space, since they are energy related. Since light-speed is inversely linked to both these properties, if the energy density of the vacuum increases, light-speed will decrease uniformly throughout the cosmos. Indeed, in 1990 Scharnhorst [51] and Barton [20] demonstrated that a lessening of the energy density of a vacuum would produce a higher velocity for light. This is explicable in terms of the QED approach. The virtual particles that make up the “*seething vacuum*” can absorb a photon of light and then re-emit it when they annihilate. This process, while fast, takes a finite time. The lower the energy density of the vacuum, the fewer virtual particles will be in the path of light photons in transit. As a consequence, the fewer absorptions and re-emissions which take place over a given distance, the faster light travels over that distance [52, 53].

However, the converse is also true. The higher the energy density of the vacuum, the more virtual particles will interact with the light photons in a given distance, and so the slower light will travel. Similarly, when light enters a transparent medium such as glass, similar absorptions and re-emissions occur, but this time it is the atoms in the glass which absorb and re-emit the light photons. This is why light slows as it travels through a denser medium. Indeed, the more closely packed the atoms, the slower light will travel as a greater number of interactions occur in a given distance. In a recent illustration of this light-speed was reduced to 17 metres/second as it passed through extremely closely packed sodium atoms near absolute zero [54]. All this is now known from experimental physics. This agrees with Barnett’s comments in *Nature* [11] that “*The vacuum is certainly a most mysterious and elusive object...The suggestion that the value of the speed of light is determined by its structure is worthy of serious investigation by theoretical physicists.*”

However, there is another way of looking at this that will also prove useful. The velocity of a wave motion such as light is proportional to the square root of the ratio of the elasticity N to the inertia K of the medium

in which it is travelling. Inertia may be defined as the ability to resist a change of velocity. From this standard relationship the following equation may be derived [55]

$$c^2 = N / K$$

Now since space has uniform properties throughout the cosmos, and is a non-dispersive medium, the wave energy associated with N and K will be equally distributed between them. This means that c is directly proportional to N, the vacuum elasticity, and proportional to 1/K, the inverse of the inertial property of space. On this new model, the redshift measurements imply that the vacuum elasticity N is declining exponentially at astronomical distances. In turn, this would also indicate that light speed 'c' is dropping exponentially as 'c' is directly proportional to N. For each redshift quantum change, the speed of light has apparently changed by a significant amount. The precise quantity is dependent upon the value adopted for the *Hubble constant* which links a galaxy's redshift with its distance.

AN OBSERVED DECLINE IN LIGHT-SPEED

The question then arises as to whether or not any other observational evidence exists that the speed of light has diminished with time. Surprisingly, some 40 articles about this very matter appeared in the scientific literature from 1926 to 1944 [56]. Some important points emerge from this literature. In 1944, despite a strong preference for the constancy of atomic quantities, N. E. Dorsey [57] was reluctantly forced to admit: "*As is well known to those acquainted with the several determinations of the velocity of light, the definitive values successively reported ... have, in general, decreased monotonously from Cornu's 300.4 megametres per second in 1874 to Anderson's 299.776 in 1940...*" Even Dorsey's own re-working of the data could not avoid that conclusion.

However, the decline in the measured value of 'c' was noticed much earlier. In 1886, Simon Newcomb reluctantly concluded that the older results obtained around 1740 were in agreement with each other, but they indicated 'c' was about 1% higher than in his own time [58], the early 1880's. In 1941 history repeated itself when Birge made a parallel statement while writing about the 'c' values obtained by Newcomb, Michelson, and others around 1880. Birge was forced to concede that "... *these older results are entirely consistent among themselves, but their average is nearly 100 km/s greater than that given by the eight more recent results*" [59]. Each of these three eminent scientists held to a belief in the absolute constancy of 'c'. This makes their careful admissions about the experimentally declining values of measured light speed more significant.

EXAMINING THE DATA

The data obtained over the last 320 years at least imply a decay in 'c' [56]. Over this period, all 163 measurements of light-speed by 16 methods reveal a non-linear decay trend. Evidence for this decay trend exists within each measurement technique as well as overall. Furthermore, an initial analysis of the behaviour of a number of other atomic constants was made in 1981 to see how they related to 'c' decay. On the basis of the measured value of these "constants", it became apparent that energy was being conserved throughout the process of 'c' variation. In all, confirmatory trends appear in 475 measurements of 11 other atomic quantities by 25 methods. Analysis of the most accurate atomic data reveals that the trend has a consistent magnitude in all the other atomic quantities that vary synchronously with light-speed [56].

All these measurements have been made during a period when there have been no quantum increases in the energy of atomic orbits. These observations reinforce the conclusion that, between any proposed quantum jumps, energy is conserved in all relevant atomic processes, as no extra energy is accessible to the atom from the ZPF. Because energy is conserved, the c-associated atomic constants vary synchronously with c, and the existing order in the cosmos is not disrupted or intruded upon. Historically, it was this very behaviour of the various constants, indicating that energy was being conserved, which was a key factor in the development of the 1987 Norman-Setterfield report, *The Atomic Constants, Light And Time* [56]. The mass of data supporting these conclusions comprises some 638 values measured by 43 methods. Montgomery and Dolphin did a further extensive statistical analysis on the data in 1993 and concluded that the results supported the 'c' decay proposition if energy was conserved [60]. The analysis was developed

further and formally presented in August 1994 by Montgomery [61]. These papers answered questions related to the statistics involved and have not yet been refuted.

ATOMIC QUANTITIES AND ENERGY CONSERVATION

Planck's constant and mass are two of the quantities which vary synchronously with 'c'. Over the period when 'c' has been measured as declining, Planck's constant 'h' has been measured as increasing as documented in the 1987 Report. The most stringent data from astronomy reveal 'hc' must be a true constant [62 - 65]. Consequently, 'h' must be proportional to '1/c' exactly. This is explicable in terms of the SED approach since, as mentioned above, 'h' is essentially a measure of the strength of the zero-point fields (ZPF). If the ZPE is increasing, so, in direct proportion, must 'h'. As noted above, an increasing ZPE also means 'c' must drop. In other words, as the energy density of the ZPF increases, 'c' decreases in such a way that 'hc' is invariant. A similar analysis could be made for other time-varying "constants" that change synchronously with 'c'.

This analysis reveals some important consequences resulting from Einstein's famous equation [$E = mc^2$], where 'E' is energy, and 'm' is mass. Data listed in the Norman/Setterfield Report confirm the analysis that 'm' is proportional to $1/c^2$ within a quantum interval, so that energy (E) is unaffected as 'c' varies. Haisch, Rueda and Puthoff independently verify that when the energy density of the ZPF decreases, mass also decreases. They confirm that 'E' in Einstein's equation remains unaffected by these synchronous changes involving 'c' [16].

If we continue this analysis, the behaviour of mass 'm' is found to be very closely related to the behaviour of the **Gravitational constant** 'G' and gravitational phenomena. In fact 'G' can be shown to vary in such a way that 'Gm' remains invariant at all times. This relationship between 'G' and 'm' is similar to the relationship between Planck's constant and the speed of light that leaves the quantity 'hc' unchanged. The quantity 'Gm' always occurs as a united entity in the relevant gravitational or orbital equations [66]. Therefore, gravitational and orbital phenomena will be unchanged by varying light speed as will planetary periods and distances [67]. In other words, acceleration due to gravity, weight, and planetary orbital years, remain independent of any variation of 'c'. As a result, astronomical orbital periods of the earth, moon, and planets form an independent time-piece, a dynamical clock, with which it is possible to compare atomic processes.

THE BEHAVIOUR OF ATOMIC CLOCKS

This comparison between dynamical and atomic clocks leads to another aspect of this discussion. Observations reveal that a higher speed of light implies that some atomic processes are proportionally faster. This includes atomic frequencies and the rate of ticking of atomic clocks. In 1934 'c' was experimentally determined to be varying, but measured wavelengths of light were experimentally shown to be unchanged. Professor Raymond T. Birge, who did not personally accept the idea that the speed of light could vary, nevertheless stated that the observational data left only one conclusion. He stated that if 'c' was actually varying and wavelengths remained unchanged, this could only mean "*the value of every atomic frequency...must be changing*" [68].

Birge was able to make this statement because of an equation linking the wavelength 'W' of light, with frequency 'F', and light-speed 'c'. The equation reads $c = FW$. If 'W' is constant and 'c' is varying, then 'F' must vary in proportion to 'c'. Furthermore, Birge knew that the frequency of light emitted from atoms is directly proportional to the frequency of the revolution of atomic particles in their orbits [42]. All atomic frequencies are therefore directly proportional to 'F', and so also directly proportional to 'c', just as Birge indicated.

The run-rate of atomic clocks is governed by atomic frequencies. It therefore follows that these clocks, in all their various forms, run at a rate proportional to c. The atomic clock is thereby c-dependent, while the orbital or dynamical clock ticks independently at a constant rate. In 1965, Kovalevsky pointed out the converse of this. He stated that if the two clock rates were different, "*then Planck's constant as well as atomic frequencies would drift*" [69]. This is precisely what the observations reveal.

This has practical consequences in the measurements of 'c'. In 1949 the frequency-dependent ammonia-quartz clock was introduced and became standard in many scientific laboratories [70]. But by 1967, atomic clocks had become uniformly adopted as timekeepers around the world. Methods that use atomic clocks to measure 'c' will always fail to detect any changes in light-speed, since their run-rate varies directly as 'c' varies. This is evidenced by the change in character of the 'c' data following the introduction of these clocks. This is why the General Conference on Weights and Measures meeting in Paris in October of 1983 declared 'c' an absolute constant [71]. Since then, any change in the speed of light would have to be inferred from measurements other than those involving atomic clocks.

COMPARING ATOMIC AND DYNAMIC CLOCKS

However, this problem with frequencies and atomic clocks can actually supply additional data to work with. It is possible in principle to obtain evidence for speed of light variation by comparing the run-rate of atomic clocks with that of dynamical clocks. When this is done, a difference in run-rate is noted. Over a number of years up to 1980, Dr. Thomas Van Flandern of the US Naval Observatory in Washington examined data from lunar laser ranging using atomic clocks, and compared their data with data from dynamical, or orbital, clocks. From this comparison of data, he concluded that "*the number of atomic seconds in a dynamical interval is becoming fewer. Presumably, if the result has any generality to it, this means that atomic phenomena are slowing down with respect to dynamical phenomena*" [72]. Van Flandern has more recently been involved in setting the parameters running the clocks in the Global Positioning System of satellites used for navigation around the world. His clock comparisons indicated that atomic phenomena were slowing against the dynamical standard until about 1980. This implies that 'c' was continuing to slow, regardless of the results obtained using the frequency-dependent measurements of recent atomic clocks.

AN OSCILLATION IS INVOLVED

These clock comparisons are useful in another way. The atomic dates of historical artifacts can be approximated via radiometric dating. These dates can then be compared with actual historical, or orbital, dates. This comparison of clocks allows us to examine the situation prior to 1678 when the Danish astronomer Roemer made the first measurement of the speed of light. When this comparison is done, light-speed behaviour is seen to include an oscillation on top of the exponential decay pattern revealed by the redshift. This evidence seems to suggest that the oscillation peaked somewhere around 500 AD. Furthermore, it is of interest to note that measurements of several atomic constants associated with 'c' also seem to indicate that the 'c' decay curve may have bottomed out around 1980 and has started to increase again. More data are needed before a positive statement can be made.

Because the oscillation is small, it only becomes apparent as the exponential curve tapers off. As both Close [73] and D'azzo & Houpis [74] pointed out in 1966, this is typical of many physical systems. The complete response of a system to an input of energy comprises two parts: the forced response and the free or natural response. This can be illustrated by a number of mechanical or electrical systems. The forced response comes from the injection of energy into the system. The free response is the system's own natural period of oscillation. The two together describe the complete behaviour of the system. In this new model, the exponential curve represents the energy injection into the system as the initial elastic tension changed its form into the ZPE, while the oscillation comes from the free response of the cosmos to this energy injection. This dual process has affected atomic behaviour and light-speed throughout the cosmos.

LIGHT-SPEED AND THE EARLY COSMOS

The issue of light-speed in the early cosmos is one which has received some attention recently in several peer-reviewed journals. Starting in December 1987, the Russian physicist V. S. Troitskii from the Radiophysical Research Institute in Gorky published a twenty-two page analysis in *Astrophysics and Space Science* regarding the problems cosmologists faced with the early universe. He looked at a possible solution if it was accepted that light-speed continuously decreased over the lifetime of the cosmos, and the

associated atomic constants varied synchronously. He suggested that, at the origin of the cosmos, light may have travelled at 10^{10} times its current speed. He concluded that the cosmos was static and not expanding.

In 1993, J. W. Moffat of the University of Toronto, Canada, had two articles published in the *International Journal of Modern Physics D* (see also [75]). He suggested that there was a high value for 'c' during the earliest moments of the formation of the cosmos, following which it rapidly dropped to its present value. Then, in January 1999, a paper in *Physical Review D* by Andreas Albrecht and Joao Magueijo, entitled "A Time Varying Speed Of Light As A Solution To Cosmological Puzzles" received a great deal of attention. These authors demonstrated that a number of serious problems facing cosmologists could be solved by a very high initial speed of light.

Like Moffat before them, Albrecht and Magueijo isolated their high initial light-speed and its proposed dramatic drop to the current speed to a very limited time during the formation of the cosmos. However, in the same issue of *Physical Review D* there appeared a paper by John D. Barrow, Professor of Mathematical Sciences at the University of Cambridge. He took this concept one step further by proposing that the speed of light has dropped from the value proposed by Albrecht and Magueijo down to its current value over the lifetime of the universe.

An article in *New Scientist* for July 24, 1999, summarised these proposals in the first sentence. "Call it heresy, but all the big cosmological problems will simply melt away, if you break one rule, says John D. Barrow – the rule that says the speed of light never varies." Interestingly, the initial speed of light proposed by Albrecht, Magueijo and Barrow is 10^{60} times its current speed. In contrast, the redshift data give a far less dramatic result. The most distant object seen in the Hubble Space Telescope has a redshift, 'z', of 14. This indicates light-speed was about 9×10^8 greater than now. At the origin of the cosmos this rises to about 2.5×10^{10} times the current value of c, more in line with Troitskii's proposal, and considerably more conservative than the Barrow, Albrecht and Magueijo estimate. This lower, more conservative estimate is also in line with the 1987 Norman-Setterfield Report.

IMPLICATIONS OF THIS PROPOSED MODEL

(1). Energy output from distant astronomical sources

Distant quasars [76] and gamma ray bursts [77] have an intense stream of redshifted photons coming from them. However, similar objects which are known to be closer, do not have the same energy output. There is a phenomena that seems to be related to distance resulting in a dilemma regarding the energy source for both types of objects.

It is normally assumed that redshifted photons outside our galaxy were emitted with the same energy as unshifted photons within our galaxy. However we know that redshifted photons have lower energy. This model accepts that these photons had a lower intrinsic energy at emission. It can be shown that the energy output by stars will remain approximately the same at all quantum jumps. Therefore stars must have emitted more lower energy photons per unit of volume in times past, thus preserving the approximate total energy output.

This explains why we see the much more intense streams of redshifted photons from distant astronomical objects as compared with similar nearby objects in order to maintain the total energy output, more lower energy photons were emitted earlier.

(2). Quantum "shells"

This model assumes each quantum change occurs instantaneously throughout the cosmos. Yet a finite time is taken for light emitted by atomic processes to reach the observer. Consequently, the observed redshift will appear to be quantised in spherical shells centred about any observer anywhere in the universe. The distance between shell boundaries will be constant because of the unique behaviour which is described by equations derived from the observational data. This distance between shell boundaries is about 138,000

light years and marks the distance between successive redshift jumps of 2.73 km/s. All objects that emit light within that shell will have the same redshift.

(3). “Missing mass” in galaxy clusters

The relative velocities of individual galaxies within clusters of galaxies are measured by their redshift. From this redshift measurement, it has been concluded that the velocities of galaxies are too high for them to remain within the cluster for the assumed age of the universe. Therefore astronomers have been looking for the “missing mass” needed to hold such clusters together by way of gravitational forces. However, if the redshift does not represent velocity, as is currently accepted, then the problem disappears. As actual relative velocities of galaxies will be small, no mass is “missing.”(Note that this does not solve the problem of the “missing mass” within spiral galaxies which is a separate issue.)

(4). A uniform microwave background

An initial very high value for light-speed means that the radiation in the very early moments of the cosmos would be rapidly homogenised by scattering processes. This means that the radiation we observe from that time will be both uniform and smooth. This is largely what is observed with the microwave background radiation coming from all parts of the sky [78]. This model therefore provides an answer to its smoothness without the necessity of secondary assumptions about matter distribution and galaxy formation that tend to be a problem for current theories.

(5). Corrections to the atomic clock

As a consequence of knowing how light-speed and atomic clocks have behaved from the redshift, atomic and radiometric clocks can now be corrected to read actual orbital time. As a result, geological eras can have a new orbital time-scale set beside them. This will necessitate a re-orientation in our current thinking on such matters.

(6). Final note

The effects of changing the vacuum energy density uniformly throughout the cosmos have been considered in this presentation. This in no way precludes the possibility that the vacuum energy density may vary on a local astronomical scale, perhaps due to energetic processes. In such cases, dramatically divergent redshifts may be expected when two neighbouring astronomical objects are compared. Arp has listed off a number of potential instances where this explanation may be valid [79, 80].

SUMMARY

This model proposes that an initial small, hot, dense, highly energetic universe underwent rapid expansion to its current size, and remained static thereafter. The vacuum potential energy in the form of an elasticity, tension, or stress, acquired from the initial expansion, became converted exponentially into the vacuum zero-point radiation. This had two results. First, there was a progressive decline in light-speed. Concurrently, atomic particle and orbital energies throughout the cosmos underwent a series of quantum increases, as more energy became available to them from the vacuum. Therefore, with increasing time, atoms emitted light that shifted in jumps towards the more energetic blue end of the spectrum. As a result, as we look back in time to progressively more distant astronomical objects, we see that process in reverse. That is to say the light of these galaxies is shifted in jumps towards the red end of the spectrum. The implications of this model solve some astronomical problems but, at the same time, challenge some current historical interpretations.

REFERENCES:

[1]. Timothy H. Boyer, “*The Classical Vacuum*”, **Scientific American**, pp.70-78, August 1985.

- [2]. Robert Matthews, “*Nothing like a Vacuum*”, **New Scientist**, p. 30-33, 25 February 1995.
- [3]. Harold E. Puthoff, “*Can The Vacuum Be Engineered For Spaceflight Applications? Overview Of Theory And Experiments*”, **NASA Breakthrough Propulsion Physics Workshop**, August 12-14, 1997, NASA Lewis Research Center, Cleveland, Ohio.
- [4]. Harold E. Puthoff, “*Everything for nothing*”, **New Scientist**, pp.36-39, 28 July 1990.
- [5]. Anonymous, “*Where does the zero-point energy come from?*”, **New Scientist**, p.14, 2 December 1989.
- [6]. Martin Harwit, “*Astrophysical Concepts*”, p. 513, Second Edition, Springer- Verlag, 1988.
- [7]. A. P. French, “*Principles of Modern Physics*”, p. 176, John Wiley & Sons, New York, 1959.
- [8]. P. W. Milonni, “*The Quantum Vacuum: An Introduction to Quantum Electrodynamics*”, Academic Press, New York, 1994.
- [9]. Timothy H. Boyer, “*Random Electrodynamics: The theory of classical electrodynamics with classical electromagnetic zero-point radiation*”, **Physical Review D**, Vol. 11:4, pp.790-808, 15 February, 1975.
- [10]. L. de la Pena, and A. M. Cetto, “*The Quantum Dice: An introduction to stochastic electrodynamics.*” Kluwer Academic Publisher, Dordrecht, 1996.
- [11]. Stephen M. Barnett, “*Photons faster than light?*”, **Nature**, Vol. 344, p. 289, 22 March, 1990.
- [12]. Kenneth W. Ford, “*Classical and Modern Physics*”, Vol. 3, p.1290, Wiley, New York, 1974.
- [13]. I. Levine et al., “*Measurement of the Electromagnetic Coupling at large momentum Transfer*”, **Physical Review Letters**, 78:3, pp. 424-427, 20 Jan 1997.
- [14]. K. Huang, “*On the Zitterbewegung of the Dirac Electron*”, **American Journal of Physics**, Vol. 20, pp.479-484, 1952.
- [15]. Bernard Haisch, Alfonso Rueda and H.E. Puthoff, “*Beyond $E = mc^2$. A First Glimpse of a Universe Without Mass*”, **The Sciences**, pp. 26-31, New York Academy of Sciences, November/December 1994.
- [16]. B. Haisch, A. Rueda, and H. E. Puthoff, “*Physics of the Zero-Point Field :Implications for Inertia, Gravitation and Mass*”, **Speculations in Science and Technology**, Vol. 20, pp. 99-114, 1997.
- [17]. E. Nelson, “*Derivation of the Schroedinger Equation from Newtonian Mechanics*”, **Physical Review**, Vol. 150, pp.1079-1085, 1966.
- [18]. Jack S. Greenberg and Walter Greiner, “*Search for the sparking of the vacuum*”, **Physics Today**, pp.24-32, August 1982.
- [19]. Walter J. Moore, “*Physical Chemistry*”, pp. 12-13, Longmans 1961.
- [20]. G. Barton, “*Faster-Than-c Light Between Parallel Mirrors*”, **Physics Letters B**, Vol. 237, No. 3,4, p. 559-562, 22 March, 1990.
- [21]. B.I. Bleaney and B. Bleaney, “*Electricity and Magnetism*”, p.242, Oxford, at the Clarendon Press, 1962.
- [22]. R. L. Lehrman and C. Swartz, “*Foundations of Physics*”, pp. 510-511, Holt, Rinehart and Winston Inc., 1969.
- [23]. H. E. Puthoff, “*Ground state of hydrogen as a zero-point-fluctuation-determined state*”, **Physical Review D**, Vol. 35, No. 10, pp. 3266-3269, 15 May, 1987.
- [24]. Donald Goldsmith, “*The Evolving Universe*”, Second Edition, pp. 108-110, Addison-Wesley, 1985.
- [25]. Paul Couderc, “*The Wider Universe*”, p. 92, Arrow Science Series, Hutchinson, London, 1960.
- [26]. William G. Tifft, “*Discrete States Of Redshift And Galaxy Dynamics I*”, **Astrophysical Journal**, Vol. 206:38-56, 15 May, 1976.
- [27]. William G. Tifft, “*Discrete States Of Redshift And Galaxy Dynamics II: Systems Of Galaxies*”, **Astrophysical Journal**, Vol. 211:31-46, 1 Jan., 1977.
- [28]. William G. Tifft, “*Discrete States Of Redshift And Galaxy Dynamics III: Abnormal Galaxies*”, **Astrophysical Journal**, 211:377-391, 15 January, 1977.
- [29]. William G. Tifft, “*The Discrete Redshift And Asymmetry In H I Profiles*”, **Astrophysical Journal**, Vol. 221:449-455, 15 April, 1978.
- [30]. William G. Tifft, “*The Absolute Solar Motion And The Discrete Redshift*”, **Astrophysical Journal**, Vol. 221:756-775, 1 May, 1978.
- [31]. William G. Tifft, “*Periodicity In The Redshift Intervals For Double Galaxies*”, **Astrophysical Journal**, Vol. 236:70-74, 15 February, 1980.
- [32]. William G. Tifft, “*Structure Within Redshift-Magnitude Bands*”, **Astrophysical Journal**, Vol. 233:799-808, 1 November, 1979.
- [33]. William G. Tifft, “*Quantum Effects In The Redshift Intervals For Double Galaxies*”, **Astrophysical Journal**, Vol. 257:442-499, 15 June, 1982.

- [34]. William G. Tifft, “*Double Galaxy Investigations II*”, **Astrophysical Journal**, Vol. 262:44-47, 1 November, 1982.
- [35]. John Gribbin, “*Galaxy red shifts come in clumps*”, **New Scientist**, pp.20-21, 20 June, 1985.
- [36]. W. J. Cocke and W. G. Tifft, “*Redshift Quantisation In Compact Groups Of Galaxies*”, **Astrophysical Journal**, Vol. 268:56-59, 1 May, 1983. Also Cocke and Tifft, **Astrophysical Journal**, Vol. 287:492. Also Cocke, **Astrophysics Letters**, Vol. 23, p. 239; **Astrophysical Journal**, Vol. 288, p.22.
- [37]. T. Beardsley, “*Quantum Dissidents*”, **Scientific American**, December 1992.
- [38]. John Gribbin, “*Riddle of the Red Shift*”, **New Scientist**, p.17, 9 July, 1994.
- [39]. R. Matthews, “*Do Galaxies Fly Through The Universe In Formation?*”, **Science**, Vol. 271:759, 1996.
- [40]. W. G. Tifft, “*Properties Of The Redshift III: Temporal Variation*”, **Astrophysical Journal**, Vol. 382:396-415, 1 December, 1991.
- [41]. J. Audouze and G. Israel, “*Cambridge Atlas of Astronomy*”, p. 382, Cambridge/Newnes, 1985.
- [42]. A. P. French, “*Principles Of Modern Physics*”, p.103-121, John Wiley & Sons, New York, 1959.
- [43]. H. E. Puthoff, “*Polarizable-Vacuum (PV) representation of general relativity*”, published by Institute For Advanced Studies at Austin, Texas, September 1999.
- [44]. John Gribbin, “*More to electrons than meets the eye*”, **New Scientist**, p. 15, 25 January, 1997.
- [45]. Robert M. Eisberg, “*Fundamentals Of Modern Physics*”, p.137, Wiley, New York, 1961.
- [46]. M. Russell Wehr and James A. Richards Jr., “*Physics Of The Atom*”, pp. 108,196, Addison-Wesley, 1960.
- [47]. Peter Fong, “*Elementary Quantum Mechanics*”, p. 16, Addison-Wesley, 1962.
- [48]. Paul S. Wesson, “*Cosmology and Geophysics*”, pp. 64-66, Adam Hilger Ltd., Bristol, 1978.
- [49]. Martin Harwit, op. cit., pp. 514-517.
- [50]. Steven Weinberg, “*The First Three Minutes*”, p. 138, Fontana Paperbacks, Flamingo Edition, 1983.
- [51]. K. Scharnhorst, “*On Propagation Of Light In The Vacuum Between Plates*”, **Physics Letters B**, Vol. 236:3, pp.354-359, 22 February, 1990.
- [52]. Marcus Chown, “*Can photons travel ‘faster than light’?*”, **New Scientist**, p. 12, 7 April, 1990.
- [53]. Anonymous, “*Secret of the vacuum: Speedier light*”, **Science News**, Vol. 137,p.303, 12 May, 1990.
- [54]. Philip F. Schewe and Ben Stein, “*Light Has Been Slowed To A Speed Of 17m/s*”, **American Institute of Physics, Bulletin of Physics News**, Number 415, 18 February, 1999.
- [55]. R. L. Lehrman and C. Swartz, “*Foundations of Physics*”, pp. 332-337, 510-511, Holt, Rinehart and Winston Inc., 1969.
- [56]. Trevor Norman and Barry Setterfield, “*Atomic Constants, Light, and Time*”, SRI International, August 1987. See detailed list under their reference [360].
- [57]. N. E. Dorsey, “*The Velocity Of Light*”, **Transactions of the American Philosophical Society**, 34, (Part 1), pp. 1-110, October, 1944.
- [58]. Simon Newcomb, “*The Velocity Of Light*”, **Nature**, pp. 29-32, 13 May, 1886.
- [59]. Raymond T. Birge, “*The General Physical Constants*”, **Reports On Progress In Physics**, Vol. 8, pp.90-101, 1941.
- [60]. Alan Montgomery and Lambert Dolphin, “*Is The Velocity Of Light Constant In Time?*”, **Galilean Electrodynamics**, Vol. 4:5, pp. 93-97, Sept./Oct. 1993.
- [61]. Alan Montgomery, “*A determination and analysis of the appropriate values of the speed of light to test the Setterfield hypothesis*”, **Proceedings of the Third International Conference on Creationism**, pp. 369-386, Creation Science Fellowship Inc., Pittsburgh, Pennsylvania, August 1994.
- [62]. J. N. Bahcall and E. E. Salpeter, “*On the interaction of radiation from distant sources with the intervening medium*”, **Astrophysical Journal**, Vol. 142,pp. 1677-1681, 1965.
- [63]. W. A. Baum and R. Florentin-Nielsen, “*Cosmological evidence against time variation of the fundamental constants*”, **Astrophysical Journal**, Vol. 209, pp. 319-329, 1976.
- [64]. J. E. Solheim et al., “*Observational evidence against a time variation in Planck’s constant*”, **Astrophysical Journal**, Vol. 209, pp. 330-334, 1976.
- [65]. P. D. Noerdlinger, “*Primordial 2.7 degree radiation as evidence against secular variation of Planck’s constant*”, **Physical Review Letters**, Vol. 30, pp.761-762, 1973.
- [66]. S. L. Martin and A. K. Connor, “*Basic Physics*”, Vol. 1, Seventh Edition, pp. 207-209, Whitcombe & Tombs, Melbourne, 1958.
- [67]. V. Canuto and S. H. Hsieh, “*Cosmological Variation Of G And The Solar Luminosity*”, **Astrophysical Journal**, Vol. 237, pp. 613-615, April 15, 1980.
- [68]. R. T. Birge, “*The Velocity Of Light*”, **Nature**, Vol. 134, pp.771-772, 1934.

- [69]. J. Kovalevsky, "Astronomical time", **Metrologia**, Vol. 1:4, pp.169-180, 1965.
- [70]. Samuel A. Goudsmit and Robert Claiborne, "Time", p. 106, Life Science Library, Time-Life International, 1967.
- [71]. T. Wilkie, "Time to Re-measure the Metre", **New Scientist**, p. 258, 27 October, 1983.
- [72]. T. C. Van Flandern, "Is the Gravitational Constant Changing?", **Precision Measurements and Fundamental Constants II**, NBS (US) Special Publication 617, B. N. Taylor and W. D. Phillips eds., pp.625-627, 1984.
- [73]. C. M. Close, "The Analysis of Linear Circuits", p. 476, Harcourt, Brace, and World Inc., 1966.
- [74]. J. J. D'azzo and C. H. Houppis, "Feedback Control System Analysis and Synthesis", pp. 257-259, McGraw Hill International Edition, 1966.
- [75]. M. A. Clayton and J. W. Moffat, "Dynamical mechanism for varying light velocity as a solution to cosmological problems", **Physics Letters B** 460:(3-4), pp.263-270, 12 August, 1999.
- [76]. J. Audouze and G. Israel, op. cit., p.358. D. Goldsmith, op. cit., pp. 182-183.
- [77]. R. Cowen, "Do supernovas generate gamma-ray bursts?", **Science News**, Vol.155, p. 263, 24 April, 1999.
- [78]. Martin Harwit, op. cit., pp. 177-180.
- [79]. Halton Arp, "Quasars, Redshifts and Controversies", Interstellar Media, Berkeley, California, 1987.
- [80]. Halton Arp, "Seeing Red: Redshifts, Cosmology and Academic Science", Apeiron, Montreal, 1998.

© BARRY SETTERFIELD - 15 November 1999 Minor edits 16 November 1999.
